

Washington

Drugged to death

Accidental overdoses from Rx cocktails alarm military officials

By Andrew Tilghman
atilghman@militarytimes.com
and Brendan McGarry
bmcgarry@militarytimes.com

At least 32 soldiers and Marines assigned to their services' most-supervised medical units for wounded troops have died of accidental prescription drug overdoses since 2007.

The 30 soldiers and two Marines overdosed while under the care of special Army Warrior Transition Units or the Marine Corps Wounded Warrior Regiment, created three years ago to tightly focus care and attention on troops suffering from severe physical and psychiatric problems as a result of combat.

Most of the troops had been prescribed "drug cocktails," combinations of drugs including pain killers, sleeping pills, antidepressants and anti-anxiety drugs, interviews and records show. In all cases, suicide was ruled out.

Army officials say the deaths are often complicated by troops mixing medications with alcohol, taking their own medications incorrectly or without a prescription.

It is unclear how many troops across the entire military have died from drug toxicity. Pentagon officials have not provided information about accidental drug deaths across the military despite a Military Times Freedom of Information Act request submitted nearly two months ago. Data on military deaths is compiled by the Armed Forces Institute of Pathology and maintained at the Pentagon's Defense Manpower Data Center.

The Army deaths have shocked that service's medical community and prompted an internal review. But despite a "safety standoff" in

January 2009, the number of fatalities continued to rise last year — to 15 in 2009, up from 11 the year before. Meanwhile, the total number of soldiers assigned to the 29 WTUs nationwide dropped from about 12,000 to about 9,000.

The internal review found the biggest risk factor may be putting a soldier on numerous drugs simultaneously, a practice known as polypharmacy. According to an Army analysis from June 2009, about 9 percent of WTU patients — 800 soldiers — were prescribed combinations of drugs including pain, psychiatric and sleep medications.

As a result, the Army medical community began questioning the practice of polypharmacy and has overhauled the way it prescribes, distributes and monitors the riskiest drugs.

An Army Medical Command memo dated May 14, 2009, highlighted the risks: "Certain prescription medications, alone or in combination, may cause adverse side effects that may prove lethal. These high-risk medications include, but are not limited to, narcotic analgesics, anxiolytics, and anti-seizure and insomnia medications."

In a handwritten note at the bottom of that memo, Army Surgeon General Lt. Gen. Eric Schoomaker added: "Closer oversight of polypharmaceutical use by our patients can be life-saving."

New rules and guidance to reduce drug toxicity deaths over the past two years include:

- Warning Army doctors to be "judicious in the use of psychoactive medications."

- Requiring soldiers to sign consent forms stating that they fully understand the potential risks



Too many prescriptions, too little talk

By Andrew Tilghman
atilghman@militarytimes.com

Army Warrant Officer 1 Judson Mount was taking several medications simultaneously while recovering from severe shrapnel wounds at the Warrior Transition Unit in San Antonio.

The painkiller Tramadol and the antidepressant Zoloft were a high-risk combination, medical experts say, and it required close supervision.

But Mount was dead of an accidental drug overdose in the WTU barracks for two days before anyone

found the married father of two.

The former enlisted tank commander who deployed to Iraq twice was found, forgotten and alone, on April 7, 2009, in his room next to several jars of pills. The cause of death was an accidental overdose of Tramadol. The "contributory effects" of the antidepressant "could not be excluded," according to the military autopsy report.

Whatever killed her son, Joyce Mount, a 63-year-old retired bank worker in Tennessee, does not blame the Army.

"It was a person — a pharmacist

or a doctor or something — not the Army," said Mount, whose father was a retired Air Force senior master sergeant. "The Army's been good to me. They've been good to all of us. They were here at the funeral. But I feel like somebody in the system, somebody has failed or messed up."

WO1 Mount was one of at least 32 service members to die from an accidental overdose of prescription drugs while under the care of what are supposed to be the military's most highly supervised medical units during the past three years.

related to the drugs.

- Prohibiting some soldiers from using more than one doctor to obtain medications.

- Reducing standard prescriptions for high-risk soldiers from 90-day supplies to a seven-day supply.

- Establishing alcohol-free zones in WTU barracks and issuing no-alcohol orders to some heavily medicated soldiers.

Robert Moore, a spokesman for

Warrior Transition Command, which oversees the WTUs, told Military Times that none of the fatalities resulted from a soldier taking his medications as instructed. Rather, they involved soldiers who took too much medication, took medication without a prescription, or mixed medication with alcohol or illegal drugs.

"These are individuals," he said. "They will make some of

their own decisions."

Moore said the rate of deaths has decreased due to the series of new safety measures. So far this year, two soldiers have died from accidental drug overdoses, and several determinations about causes of death are pending, according to interviews.

Nevertheless, the problem has become a priority for the Army's top leaders.



SAMUEL M. SIMPKINS/
THE TENNESSEAN

Joyce Mount's son Judd died of an accidental drug overdose at a Warrior Transition Unit in San Antonio.

Army Sgt. Franklin Barnett, a 29-year-old combat engineer and father of three, also died while under the care of the WTU in San Antonio. He was wounded by a car bomb in Iraq and received a Purple Heart in October 2008. His widow blamed his June 2009 death on communication failures by Army doctors.

"If the doctors would talk to each other, then they wouldn't have a problem," Diane Barnett said. "He was on four different kinds of medication that pretty much clashed with each other."

"With two drug-related deaths thus far this year, we are not content that we are solving this problem and continue to look at every possible avenue to further reduce the risk of such events, not only in the WTUs but across our force," Army Vice Chief of Staff Gen. Peter Chiarelli told Military Times earlier this month.

The military has a computer system designed to warn doctors

Franklin Barnett was taking "antidepressants and sleeping pills," his wife said — adding that he may not have taken his medications as prescribed. "He was forgetful — he probably forgot that he took his med and he took some more."

Accidental drug overdoses in the Army WTUs began to draw public attention nearly three years ago after the death of Sgt. Gerald Cassidy, found dead in his barracks at Fort Knox, Ky., in September 2007.

He died after taking a mix of drugs that included several strong narcotic painkillers and Celexa, an antidepressant. His military autopsy concluded that the drugs' "combined synergistic" effects caused cardiac arrest.

Sen. Evan Bayh, D-Ind., heard about the death and, as a member of the Senate Armed Services Committee, demanded an Army investigation.

"The enemy could not kill him, but our own government did," Bayh said in November 2007 during a committee hearing.

Then-Army Secretary Pete Geren called the circumstances of Cassidy's death "unacceptable," and Army leaders promised to investigate.

Some family members remain angry at the Army.

"They still haven't owned up to it and said, 'You know what? We killed your husband,'" said Susan Nichols, widow of Sgt. Robert Nichols, 32, who died at the WTU in San Antonio.

Diagnosed with post-traumatic stress disorder and suffering primarily from psychiatric problems, Robert Nichols was taking a mix of 11 drugs that left him groggy and confused during the last few weeks of his life. They included Percocet, Valium, Celexa, the antipsychotic Seroquel, and Depakote, an anti-seizure drug used to treat major depression and bipolar disorder, Susan said.

"I blame those who prescribed the pills and were watching over him," she said. "They should have been able to see the signs that something was wrong." □

when individuals receive drugs that may cause adverse reactions. But doctors are able to easily override the warning notification and allow patients to receive high-risk combinations, military records show.

The details underlying each death are unique. Army Sgt. Gerald Cassidy died in 2007 after writing in his journal that he was unsure how much methadone he

SPIKE IN DRUG ORDERS

Defense Logistics Agency data show a steep increase in prescription orders for certain medications from 2001 to 2009:

Anti-anxiety/sedatives (Includes anxiolytics and insomnia medications) **170%**

Anticonvulsants (Includes anti-seizure medications) **68**

Antidepressants **39**

Antipsychotics **234**

Painkillers (Includes narcotic analgesics) **193**

Source: DLA BRYAN SMITH/STAFF

had taken, his family said.

Army Warrant Officer 1 Judson Mount died in April 2009 after trying a new, higher-dosage patch that releases the narcotic painkiller fentanyl, his mother said.

And Spc. Franklin Barnett died in June 2009 shortly after spending a weekend with his wife and children and appearing to be in good health, his wife said.

Unlike casualties in Iraq or Afghanistan, these fatalities can be avoided through better management of the health care units, said Col. (Dr.) Steven Swann, command surgeon for the Warrior Transition Command.

"Losing a soldier in combat is an expected and understood cost of ... war. But these should be preventable," Swann said. "We will do everything we can ... — more policies, more programs, more controls — to prevent every single one of these."

Meds on the rise

During the past decade — for nearly all of which the U.S. has been at war on two fronts — the military community has seen a dramatic rise in the use of the types of medications linked to the WTU deaths. For example, the military health care system's prescription orders for painkillers nearly tripled, while those for anti-seizure medications rose 68 percent, according to a recent Military Times analysis of Defense Logistics Agency data.

Many of those drugs have a similar fundamental effect on the body, slowing the central nervous system and increasing the risk that a patient's heart or breathing will stop during sleep.

"Using alcohol and illicit drugs

in combination with high-risk medications increases the potential for adverse events and death," the April 2009 Army Medical Command memo said.

The spate of deaths fuels criticism that the military medical community — and the American medical community at large — puts too much emphasis on pharmaceutical products rather than other forms of treatment.

"There is a direct correlation in the increase of use of these medications and these sudden deaths," said Dr. Bart Billings, a retired Army colonel and psychologist in San Diego who treats troubled troops and has testified before Congress about the risks linked to prescription drugs. "These are healthy young people who are dying in their sleep because some physician prescribed a combination of medications that killed them."

Many such drugs are tested and approved for use individually, but research on combinations is limited. "These medications were not tested in combination with other medications," Billings said. "They were tested only on what they would do on their own."

Billings believes the safest and most effective treatment includes various forms of talk therapy in which troops forge personal relationships with counselors while trying to identify, understand and deal with their mental health problems.

But some military doctors caution against blaming drug use in general and note that most people respond well to painkillers and psychiatric medications.

"The reasons we use these drugs is because they work," Swann said. "They are effective at managing people's pain and managing their depression."

Marine drug deaths

The Marine Corps has wrestled with similar problems.

"Medication risk management is one of the recurring hot-button topics," said Navy Capt. William Tanner, the head doctor for the Marine Corps' Wounded Warrior Regiment.

Last year, a spate of drug thefts in the barracks at Camp Lejeune, N.C., prompted the Corps to give Marines a lockbox to secure prescription drugs, Tanner said.

Some Marines with traumatic brain injuries receive personal digital assistants to help them keep track of their daily drugs.

The Corps also is developing a program that brings doctors, case-workers and Marine officials together once a week to discuss

each patient and their medications.

"We don't have a great treatment for PTSD [post-traumatic stress disorder]," Tanner said. "There are studies and recommended treatments, but none of them are great. It's hard to tell a doctor what to do. He's going to do what he thinks is best for the patient, regardless of what the guidelines say."

Suicide semblance

An accidental drug overdose initially can be confused with suicide. After Sgt. Robert Nichols died at the WTU at Fort Sam Houston, Texas, in 2008, the Army Criminal Investigation Command grilled his wife for possible evidence that his death was self-inflicted.

"The CID guys were like, 'Well, you know, was there anything that was on his plate that was too much to handle? Was there anything bothering him?'" said Susan Nichols, who now lives in Dallas. "You didn't have to be Albert Einstein to see where they were going with that. I thought, are you really trying to suggest this? This man? No."

Nichols, who deployed to Iraq in 2007 to a base south of Baghdad, sustained a traumatic brain injury after a mortar round landed near him, his wife said.

An investigation later concluded that Nichols' death was an accident. Medical records show he was taking a cocktail of 11 drugs, including Percocet, Valium, the antidepressant Celexa, the antipsychotic Seroquel, and Depakote, an anti-seizure drug used to treat major depression and bipolar disorder, his wife said.

Some psychiatric medications in the accidental overdoses come with warnings about increased risks for suicidal thoughts and actions.

The Army estimates that about 5 percent of suicides involve prescription drugs, documents show.

When the cause of death is unclear, the military can consult a forensic psychiatrist, who examines in detail the victim's life and activities and apparent frame of mind in the hours before the death. Law enforcement investigators can also be involved.

But final determinations are not always clear-cut, said Army Col. David Benedek, who teaches psychiatry at the Uniformed Services University of the Health Sciences, a Defense Department school in Bethesda, Md.

Accidents and suicides, he said, "are difficult distinctions to make sometimes, particularly if someone doesn't leave a note or indicate in any way that they were contemplating suicide." □